Quality ID #226 (NQF 0028): Preventive Care and Screening: Tobacco Use: Screening and Cessation Intervention

- National Quality Strategy Domain: Community/Population Health
- Meaningful Measure Area: Prevention and Treatment of Opioid and Substance Use Disorders

#### **2021 COLLECTION TYPE:**

MIPS CLINICAL QUALITY MEASURES (CQMS)

#### **MEASURE TYPE:**

**Process** 

#### **DESCRIPTION:**

Percentage of patients aged 18 years and older who were screened for tobacco use one or more times within 12 months **AND** who received tobacco cessation intervention if identified as a tobacco user

#### **INSTRUCTIONS:**

This measure is to be submitted a minimum of <u>once per performance period</u> for patients seen during the performance period. This measure is intended to reflect the quality of services provided for preventive screening for tobacco use. This measure may be submitted by Merit-based Incentive Payment System (MIPS) eligible clinicians who provided the measure-specific denominator coding. For this implementation of the measure, the 12 month look back period includes the program year and the year prior. For Quality Payment Program (QPP) 2021, the 12 month period would be from 1/1/2021-12/31/2021.

#### This measure will be calculated with 3 performance rates:

- 1) Percentage of patients aged 18 years and older who were screened for tobacco use one or more times within 12 months
- 2) Percentage of patients aged 18 years and older who were identified as a tobacco user who received tobacco cessation intervention
- 3) Percentage of patients aged 18 years and older who were screened for tobacco use one or more times within 12 months AND who received tobacco cessation intervention if identified as a tobacco user

The denominator of submission criteria 2 is a subset of the resulting numerator for submission criteria 1, as submission criteria 2 is limited to assessing if patients identified as tobacco users received an appropriate tobacco cessation intervention. For all patients, submission criteria 1 and 3 are applicable, but submission criteria 2 will only be applicable for those patients who are identified as tobacco users. Therefore, data for every patient that meets the age and encounter requirements will only be submitted for submission criteria 1 and 3, whereas data submitted for submission criteria 2 will be for a subset of patients who meet the age and encounter requirements, as the denominator has been further limited to those who were identified as tobacco users.

**NOTE:** Patient encounters for this measure conducted via telehealth (e.g., encounters coded with GQ, GT, 95, or POS 02 modifiers) are allowable.

#### **Measure Submission Type:**

Measure data may be submitted by individual MIPS eligible clinicians, groups, or third party intermediaries. The listed denominator criteria are used to identify the intended patient population. The numerator options included in this specification are used to submit the quality actions as allowed by the measure. The quality-data codes listed do not need to be submitted by MIPS eligible clinicians, groups, or third party intermediaries that utilize this modality for submissions; however, these codes may be submitted for those third party intermediaries that utilize Medicare Part B claims data. For more information regarding Application Programming Interface (API), please refer to the Quality Payment Program (QPP) website.

#### THERE ARE THREE SUBMISSION CRITERIA FOR THIS MEASURE:

1) All patients who were screened for tobacco use

#### AND

2) All patients who were identified as a tobacco user and who received tobacco cessation intervention

#### AND

3) All patients who were screened for tobacco use and, if identified as a tobacco user received tobacco cessation intervention, or identified as a tobacco non-user

This measure contains three submission criteria which aim to identify patients who were screened for tobacco use (submission criteria 1), patients who were identified as tobacco users and who received tobacco cessation intervention (submission criteria 2), and a comprehensive look at the overall performance on tobacco screening and cessation intervention (submission criteria 3). By separating this measure into various submission criteria, the MIPS eligible professional or MIPS eligible clinician will be able to better ascertain where gaps in performance exist, and identify opportunities for improvement. The overall rate (submission criteria 3) can be utilized to compare performance to published versions of this measure prior to the 2018 performance year, when the measure had a single performance rate. For accountability reporting in the CMS MIPS program, the rate for submission criteria 2 is used for performance.

#### SUBMISSION CRITERIA 1: ALL PATIENTS WHO WERE SCREENED FOR TOBACCO USE

#### **DENOMINATOR (SUBMISSION CRITERIA 1):**

All patients aged 18 years and older seen for at least two visits or at least one preventive visit during the measurement period

**DENOMINATOR NOTE:** \*Signifies that this CPT Category I code is a non-covered service under the Medicare Part B Physician Fee Schedule (PFS). These non-covered services should be counted in the denominator population for MIPS CQMs.

#### <u>Denominator Criteria (Eligible Cases):</u>

Patients aged ≥ 18 years

#### AND

**At least two patient encounters during the performance period (CPT):** 90791, 90792, 90832, 90834, 90837, 90845, 92002, 92004, 92012, 92014, 92521, 92522, 92523, 92524, 92540, 92557, 92625, 96156, 96158, 97161, 97162, 97163, 97165, 97166, 97167, 97168, 99202, 99203, 99204, 99205, 99212, 99213, 99214, 99215, 99341, 99342, 99343, 99344, 99345, 99347, 99348, 99349, 99350 **OR** 

At least one preventive encounter during the performance period (CPT or HCPCS): 99385\*, 99386\*, 99387\*, 99395\*, 99396\*, 99397\*, 99401\*, 99402\*, 99403\*, 99404\*, 99411\*, 99412\*, 99429\*, G0438, G0439

#### **NUMERATOR (SUBMISSION CRITERIA 1):**

Patients who were screened for tobacco use at least once within 12 months

#### **Definitions:**

**Tobacco Use** – Includes any type of tobacco.

**NUMERATOR NOTE:** To satisfy the intent of this measure, a patient must have at least one tobacco use screening during the 12-month period. If a patient has multiple tobacco use screenings during the 12-month period, only the most recent screening, which has a documented status of tobacco user or tobacco non-user, will be used to satisfy the measure requirements.

In the event that a patient is screened for tobacco use and tobacco status is unknown, submit G9905. Denominator Exception(s) are determined on the date of the most recent denominator eligible encounter for all submission criteria.

**Numerator Options:** 

Performance Met: Patient screened for tobacco use AND identified as a

tobacco user (G9902)

OR

Performance Met: Patient screened for tobacco use AND identified as a

tobacco non-user (G9903)

OR

**Denominator Exception:** Documentation of medical reason(s) for not screening

for tobacco use (e.g., limited life expectancy, other

medical reason) (G9904)

OR

Performance Not Met: Patient not screened for tobacco use, reason not

given (**G9905**)

#### SUBMISSION CRITERIA 2: ALL PATIENTS WHO WERE IDENTIFIED AS A TOBACCO USER AND WHO RECEIVED TOBACCO CESSATION INTERVENTION

#### **DENOMINATOR (SUBMISSION CRITERIA 2):**

All patients aged 18 years and older seen for at least two visits or at least one preventive visit during the measurement period who were screened for tobacco use and identified as a tobacco user

**DENOMINATOR NOTE:** \*Signifies that this CPT Category I code is a non-covered service under the Medicare Part B PFS. These non-covered services should be counted in the denominator population for MIPS CQMs.

#### **Denominator Criteria (Eligible Cases):**

Patients aged ≥ 18 years

All eligible instances when G9902 is submitted for Performance Met (patient screened for tobacco use and identified as a tobacco user) in the numerator of Submission Criteria 1

#### AND

OR

At least two patient encounters during the performance period (CPT): 90791, 90792, 90832, 90834, 90837, 90845, 92002, 92004, 92012, 92014, 92521, 92522, 92523, 92524, 92540, 92557, 92625, 96156, 96158, 97161, 97162, 97163, 97165, 97166, 97167, 97168, 99202, 99203, 99204, 99205, 99212, 99213, 99214, 99215, 99341, 99342, 99343, 99344, 99345, 99347, 99348, 99349, 99350

At least one preventive encounter during the performance period (CPT or HCPCS): 99385\*, 99386\*, 99387\*, 99395\*, 99396\*, 99397\*, 99401\*, 99402\*, 99403\*, 99404\*, 99411\*, 99412\*, 99429\*, G0438, G0439

#### **NUMERATOR (SUBMISSION CRITERIA 2):**

Patients who received tobacco cessation intervention

#### **Definitions:**

**Tobacco Cessation Intervention** Includes brief counseling (3 minutes or less), and/or pharmacotherapy. Note: For the purpose of this measure, brief counseling (e.g., minimal and intensive advice/counseling interventions conducted both in person and over the phone) qualifies for the numerator. Written self-help materials (e.g., brochures, pamphlets) and complementary/alternative therapies do not qualify for the numerator. Brief counseling also may be of longer duration or be performed more frequently, as evidence shows there is a dose-response relationship between the intensity of counseling provided (either length or frequency) and tobacco cessation rates (U.S. Preventive Services Task Force, 2015).

**NUMERATOR NOTE:** If a patient uses any type of tobacco (i.e., smokes or uses smokeless tobacco), the expectation is that they should receive tobacco cessation intervention: either counseling and/or pharmacotherapy.

This measure defines tobacco cessation counseling as lasting 3 minutes or less. Services typically provided under CPT codes 99406 and 99407 satisfy the requirement of tobacco cessation intervention, as these services provide tobacco cessation counseling for 3-10 minutes. If a patient received these types of services, submit G-code G9906.

Denominator Exception(s) are determined on the date of the most recent denominator eligible encounter for all submission criteria.

**Numerator Options:** 

Performance Met: Patient identified as a tobacco user received tobacco

cessation intervention (counseling and/or

pharmacotherapy) (G9906)

OR

**Denominator Exception:** Documentation of medical reason(s) for not providing

tobacco cessation intervention (e.g., limited life expectancy, other medical reason) (G9907)

OR

Performance Not Met: Patient identified as tobacco user did not receive

tobacco cessation intervention (counseling and/or pharmacotherapy), reason not given (**G9908**)

## SUBMISSION CRITERIA 3: ALL PATIENTS WHO WERE SCREENED FOR TOBACCO USE AND, IF IDENTIFIED AS A TOBACCO USER RECEIVED TOBACCO CESSATION INTERVENTION, OR IDENTIFIED AS A TOBACCO NON-USER

#### **DENOMINATOR (SUBMISSION CRITERIA 3):**

All patients aged 18 years and older seen for at least two visits or at least one preventive visit during the measurement period

**DENOMINATOR NOTE:** \*Signifies that this CPT Category I code is a non-covered service under the Medicare Part B PFS. These non-covered services should be counted in the denominator population for MIPS CQMs.

#### **Denominator Criteria (Eligible Cases):**

Patients aged ≥ 18 years

AND

**At least two patient encounters during the performance period (CPT):** 90791, 90792, 90832, 90834, 90837, 90845, 92002, 92004, 92012, 92014, 92521, 92522, 92523, 92524, 92540, 92557, 92625, 96156, 96158, 97161, 97162, 97163, 97165, 97166, 97167, 97168, 99202, 99203, 99204, 99205, 99212, 99213, 99214, 99215, 99341, 99342, 99343, 99344, 99345, 99347, 99348, 99349, 99350

#### OR

**At least one preventive encounter during the performance period (CPT or HCPCS):** 99385\*, 99386\*, 99387\*, 99395\*, 99396\*, 99397\*, 99401\*, 99402\*, 99403\*, 99404\*, 99411\*, 99412\*, 99429\*, G0438, G0439

#### **NUMERATOR (SUBMISSION CRITERIA 3):**

Patients who were screened for tobacco use at least once within 12 months <u>AND</u> who received tobacco cessation intervention if identified as a tobacco user

#### **Definitions:**

**Tobacco Use** – Includes any type of tobacco.

**Tobacco Cessation Intervention** – Includes brief counseling (3 minutes or less), and/or pharmacotherapy. Note: For the purpose of this measure, brief counseling (e.g., minimal and intensive advice/counseling interventions conducted both in person and over the phone) qualifies for the numerator. Written self-help materials (e.g., brochures, pamphlets) and complementary/alternative therapies do not qualify for the numerator. Brief counseling also may be of longer duration or be performed more frequently, as evidence shows there is a dose-response relationship between the intensity of counseling provided (either length or frequency) and tobacco cessation rates (U.S. Preventive Services Task Force, 2015).

**NUMERATOR NOTE:** To satisfy the intent of this measure, a patient must have at least one tobacco use screening during the 12-month period. If a patient has multiple tobacco use screenings during the 12-month period, only the most recent screening, which has a documented status of tobacco user or tobacco non-user, will be used to satisfy the measure requirements.

In the event that a patient is screened for tobacco use and identified as a user but did not receive tobacco cessation intervention or if tobacco status is unknown, submit 4004F with 8P.

If a patient uses any type of tobacco (i.e., smokes or uses smokeless tobacco), the expectation is that they should receive tobacco cessation intervention: either counseling and/or pharmacotherapy.

This measure defines tobacco cessation counseling as lasting 3 minutes or less. Services typically provided under CPT codes 99406 and 99407 satisfy the requirement of tobacco cessation intervention, as these services provide tobacco cessation counseling for 3-10 minutes. If a patient received these types of services, submit CPT II 4004F.

Denominator Exception(s) are determined on the date of the most recent denominator eligible encounter for all submission criteria.

**Numerator Options:** 

Performance Met: Patient screened for tobacco use AND received

tobacco cessation intervention (counseling,

pharmacotherapy, or both), if identified as a tobacco

user (4004F)

<u>OR</u>

Performance Met: Current tobacco non-user (1036F)

<u>OR</u>

**Denominator Exception:** Documentation of medical reason(s) for not screening

for tobacco use (e.g., limited life expectancy, other

medical reason) (4004F with 1P)

<u>OR</u>

**Denominator Exception:** Documentation of medical reason(s) for not providing

tobacco cessation intervention if identified as a tobacco user (e.g., limited life expectancy, other

Version 5.0 November 2020 CPT only copyright 2020 American Medical Association. All rights reserved.

Page 5 of 18

OR

Performance Not Met:

Tobacco screening not performed OR tobacco cessation intervention not provided, reason not otherwise specified (4004F with 8P)

#### RATIONALE:

This measure is intended to promote adult tobacco screening and tobacco cessation interventions for those who use tobacco products. There is good evidence that tobacco screening and brief cessation intervention (including counseling and/or pharmacotherapy) is successful in helping tobacco users quit. Tobacco users who are able to stop using tobacco lower their risk for heart disease, lung disease, and stroke.

#### **CLINICAL RECOMMENDATION STATEMENTS:**

The USPSTF recommends that clinicians ask all adults about tobacco use, advise them to stop using tobacco, and provide behavioral interventions and U.S. Food and Drug Administration (FDA)-approved pharmacotherapy for cessation to adults who use tobacco (Grade A Recommendation) (U.S. Preventive Services Task Force, 2015).

The USPSTF recommends that clinicians ask all pregnant women about tobacco use, advise them to stop using tobacco, and provide behavioral interventions for cessation to pregnant women who use tobacco (Grade A Recommendation) (U.S. Preventive Services Task Force, 2015).

The USPSTF concludes that the current evidence is insufficient to recommend electronic nicotine delivery systems for tobacco cessation in adults, including pregnant women. The USPSTF recommends that clinicians direct patients who smoke tobacco to other cessation interventions with established effectiveness and safety (previously stated) (Grade I Statement) (U.S. Preventive Services Task Force, 2015).

#### **COPYRIGHT:**

This Physician Performance Measure (Measure) and related data specifications are owned and copyrighted by the National Committee for Quality Assurance (NCQA). NCQA is not responsible for any use of the Measure. The Measure is not a clinical guideline and does not establish a standard of medical care and has not been tested for all potential applications.

#### THE MEASURE AND SPECIFICATIONS ARE PROVIDED "AS IS" WITHOUT WARRANTY OF ANY KIND.

NCQA makes no representations, warranties, or endorsement about the quality of any organization or physician that uses or reports performance measures and NCQA has no liability to anyone who relies on such measures or specifications.

The Measure can be reproduced and distributed, without modification, for noncommercial purposes (e.g., use by healthcare providers in connection with their practices) without obtaining approval from NCQA. Commercial use is defined as the sale, licensing, or distribution of the Measure for commercial gain, or incorporation of the Measure into a product or service that is sold, licensed or distributed for commercial gain. All commercial uses or requests for modification must be approved by NCQA and are subject to a license at the discretion of NCQA. The PCPI's and AMA's significant past efforts and contributions to the development and updating of the measure are acknowledged.

©2012-2020 National Committee for Quality Assurance. All Rights Reserved.


Limited proprietary coding is contained in the Measure specifications for user convenience. Users of proprietary code sets should obtain all necessary licenses from the owners of the code sets. NCQA disclaims all liability for use or accuracy of any third party codes contained in the specifications.

CPT® contained in the Measure specifications is copyright 2004-2020 American Medical Association. LOINC® copyright 2004-2020 Regenstrief Institute, Inc. This material contains SNOMED Clinical Terms® (SNOMED CT®)

copyright 2004-2020 International Health Terminology Standards Development Organisation. ICD-10 copyright 2020 World Health Organization. All Rights Reserved.

## 2021 Clinical Quality Measure for Quality ID #226 (NQF 0028): Preventive Care and Screening: Tobacco Use: Screening and Cessation Intervention Submission Criteria One

Disclaimer: Refer to the measure specification for the specific coding and instructions to submit this measure.




<sup>\*</sup>See the posted measure specification for specific coding and instructions to submit this measure

NOTE: Submission Frequency: Patient-Process

CPT only copyright 2020 American Medical Association. All rights reserved. The measure diagrams were developed by CMS as a supplemental resource to be used in conjunction with the measure specifications. They should not be used alone or as a substitution for the measure specification.

<sup>\*\*</sup>In the event that the tobacco status is unknown submit G9905.

#### **Submission Criteria Two**



## SAMPLE CALCULATIONS: SUBMISSION CRITERIA TWO Data Completeness= Performance Met (a³=20 patients) + Denominator Exception (b²=10 patients) + Performance Not Met (c²=10 patients) = 40 patients = 80.00% Eligible Population / Denominator (d²=50 patients) = 50 patients Performance Rate= Performance Met (a³=20 patients) = 20 patients = 66.67% Data Completeness Numerator (40 patients) – Denominator Exception (b²=10 patients) = 30 patients


NOTE: Submission Frequency: Patient-Process

CPT only copyright 2020 American Medical Association. All rights reserved. The measure diagrams were developed by CMS as a supplemental resource to be used in conjunction with the measure specifications. They should not be used alone or as a substitution for the measure specification.

<sup>\*</sup>See the posted measure specification for specific coding and instructions to submit this measure.

<sup>\*\*</sup>This measure defines tobacco cessation counseling as lasting 3 minutes or less. Services typically provided under CPT codes 99406 and 99407 satisfy the requirement of tobacco cessation intervention, as these services provide tobacco cessation counseling for 3-10 minutes. If a patient received these types of services, submit G-code G9906.

#### **Submission Criteria Three**



# SAMPLE CALCULATIONS: SUBMISSION CRITERIA THREE Data Completeness= Performance Met (a<sup>4</sup>+a<sup>5</sup>=40 patients) + Denominator Exception (b<sup>3</sup>+b<sup>4</sup>=20 patients) + Performance Not Met (c<sup>3</sup>=20 patients) = 80 patients = 80.00% Eligible Population / Denominator (d<sup>3</sup>=100 patients) = 40 patients = 66.67% Performance Rate= Performance Met (a<sup>4</sup>+a<sup>5</sup>=40 patients) = 40 patients = 66.67% Data Completeness Numerator (80 patients) - Denominator Exception (b<sup>3</sup>+b<sup>4</sup>=20 patients) = 60 patients

CPT only copyright 2020 American Medical Association. All rights reserved. The measure diagrams were developed by CMS as a supplemental resource to be used in conjunction with the measure specifications. They should not be used alone or as a substitution for the measure specification.

<sup>\*</sup>See the posted measure specification for specific coding and instructions to submit this measure.

<sup>\*\*</sup>This measure defines tobacco cessation counseling as lasting 3 minutes or less. Services typically provided under CPT codes 99406 and 99407 satisfy the requirement of tobacco cessation intervention, as these services provide tobacco cessation counseling for 3-10 minutes. If a patient received these types of services, submit 4004F.

<sup>\*\*\*</sup>In the event that a patient is identified as a user but did not receive tobacco cessation intervention submit 4004F with 8P. NOTE: Submission Frequency: Patient-Process

### 2021 Clinical Quality Measure Flow Narrative for Quality ID #226 (NQF 0028): Preventive Care and Screening: Tobacco Use: Screening and Cessation Intervention

**Disclaimer:** Refer to the measure specification for specific coding and instructions to submit this measure.

#### **Multiple Performance Rates**

#### **Submission Criteria One:**

- 1. Start with Denominator
- 2. Check Patient aged greater than or equal to 18 years:
  - a. If *Patient aged greater than or equal to 18 years* equals No, do not include in *Eligible Population/Denominator*. Stop processing.
  - b. If Patient aged greater than or equal to 18 years equals Yes, proceed to check At least two patient encounters during the performance period as listed in the Denominator\*.
- 3. Check At least two patient encounters during the performance period as listed in the Denominator\*:
  - a. If At least two patient encounters during the performance period as listed in the Denominator\* equals No, proceed to check At least one preventive encounter during the performance period as listed in the Denominator\*.
  - b. If At least two patient encounters during the performance period as listed in the Denominator\* equals Yes, include in Eligible Population/Denominator.
- 4. Check At least one preventive encounter during the performance period as listed in the Denominator\*:
  - a. If At least one preventive encounter during the performance period as listed in the Denominator\* equals No, do not include in Eligible Population/Denominator. Stop processing.
  - b. If At least one preventive encounter during the performance period as listed in the Denominator\* equals Yes, include in Eligible Population/Denominator.
- 5. Denominator Population:
  - a. Denominator Population is all Eligible Patients in the Denominator. Denominator is represented as Denominator in the Sample Calculation listed at the end of this document. Letter d<sup>1</sup> equals 100 patients in the Sample Calculation.
- 6. Start Numerator
- 7. Check Patient screened for tobacco use AND identified as a tobacco user.
  - a. If Patient screened for tobacco use AND identified as a tobacco user equals Yes, include in Data Completeness Met and Performance Met.
    - Data Completeness Met and Performance Met letter is represented in the Data Completeness and Performance Rate in the Sample Calculation listed at the end of this document. Letter a<sup>1</sup> equals 50 patients in the Sample Calculation.
  - b. If Patient screened for tobacco use AND identified as a tobacco user equals No, proceed to check Patient screened for tobacco use AND identified as a tobacco non-user.

- 8. Check Patient screened for tobacco use AND identified as a tobacco non-user.
  - a. If Patient screened for tobacco use AND identified as a tobacco non-user equals Yes, include in Data Completeness Met and Performance Met.
    - Data Completeness Met and Performance Met letter is represented in the Data Completeness and Performance Rate in the Sample Calculation listed at the end of this document. Letter a<sup>2</sup> equals 20 patients in the Sample Calculation.
  - b. If Patient screened for tobacco use AND identified as a tobacco non-user equals No, proceed to check Documentation of medical reason(s) for not screening for tobacco use.
- 9. Check Documentation of medical reason(s) for not screening for tobacco use:
  - a. If Documentation of medical reason(s) for not screening for tobacco use equals Yes, include in Data Completeness Met and Denominator Exception.
    - Data Completeness Met and Denominator Exception letter is represented in the Data Completeness and Performance Rate in the Sample Calculation listed at the end of this document. Letter b<sup>1</sup> equals 10 patients in the Sample Calculation.
  - b. If Documentation of medical reason(s) for not screening for tobacco use equals No, proceed to check Patient not screened for tobacco use, reason not given.
- 10. Check Patient not screened for tobacco use, reason not given:
  - a. If Patient not screened for tobacco use, reason not given equals Yes, include in the Data Completeness Met and Performance Not Met\*\*.
    - Data Completeness Met and Performance Not Met\*\* letter is represented in the Data
       Completeness in the Sample Calculation listed at the end of this document. Letter c<sup>1</sup> equals 10 patients in the Sample Calculation.
  - b. If Patient not screened for tobacco use, reason not given equals No, proceed to check Data Completeness Not Met.
- 11. Check Data Completeness Not Met:
  - a. If *Data Completeness Not Met*, the Quality Data Code or equivalent was not submitted. 10 patients have been subtracted from the Data Completeness Numerator in the Sample Calculation.

#### Sample Calculations: Submission Criteria One

Data Completeness equals Performance Met (a¹ plus a² equals 70 patients) plus Denominator Exception (b¹ equals 10 patients) plus Performance Not Met (c¹ equals 10 patients) divided by Eligible Population/Denominator (d¹ equals 100 patients). All equals 90 patients divided by 100 patients. All equals 90.00 percent.

Performance Rate equals Performance Met (a<sup>1</sup> plus a<sup>2</sup> equals 70 patients) divided by Data Completeness Numerator (90 patients) minus Denominator Exception (b<sup>1</sup> equals 10 patients). All equals 70 patients divided by 80 patients. All equals 87.50 percent.

\*See the posted measure specification for specific coding and instructions to submit this measure.

\*\*In the event that the tobacco status is unknown submit G9905.

NOTE: Submission Frequency: Patient-Process

The measure diagrams were developed by CMS as a supplemental resource to be used in conjunction with the measure specifications. They should not be used alone or as a substitution for the measure specification.

#### **Submission Criteria Two:**

- Start with Denominator
- 2. Check Patients aged greater than or equal to 18 years:
  - a. If Patients aged greater than or equal to 18 years equals No, do not include in Eligible Population/Denominator. Stop processing.
  - b. If Patients aged greater than or equal to 18 years equals Yes, proceed to check All eligible instances when G9902 is submitted for Performance Met (patient screened for tobacco use and identified as tobacco user) in the numerator of Submission Criteria 1.
- 3. Check All eligible instances when G9902 is submitted for Performance Met (patient screened for tobacco use and identified as tobacco user) in the numerator of Submission Criteria 1:
  - a. If All eligible instances when G9902 is submitted for Performance Met (patient screened for tobacco use and identified as tobacco user) in the numerator of Submission Criteria 1 equals No, do not include in Eligible Population/Denominator. Stop processing.
  - b. If All eligible instances when G9902 is submitted for Performance Met (patient screened for tobacco use and identified as tobacco user) in the numerator of Submission Criteria 1 equals Yes, proceed to check At least two patient encounters during the performance period as listed in Denominator\*.
- 4. Check At least two patient encounters during the performance period as listed in Denominator\*:
  - a. If At least two patient encounters during the performance period as listed in Denominator\* equals No, proceed to check At least one preventive encounter during the performance period as listed in Denominator\*.
  - b. If At least two patient encounters during the performance period as listed in Denominator\* equals Yes, include in Eligible Population/Denominator.
- 5. Check At least one preventive encounter during the performance period as listed in Denominator\*:
  - a. If At least one preventive encounter during the performance period as listed in Denominator\* equals No, do not include in Eligible Population/Denominator. Stop processing.
  - b. If At least one preventive encounter during the performance period as listed in Denominator\* equals Yes, include in Eligible Population/Denominator.
- 6. Denominator Population:
  - a. Denominator Population is all Eligible Patients in the Denominator. Denominator is represented as Denominator in the Sample Calculation listed at the end of this document. Letter d<sup>2</sup> equals 50 patients in the Sample Calculation.

- 7. Start Numerator
- 8. Check Patient identified as a tobacco user received tobacco cessation intervention (counseling and/or pharmacotherapy):
  - a. If Patient identified as a tobacco user received tobacco cessation intervention (counseling and/or pharmacotherapy) equals Yes, include in Data Completeness Met and Performance Met\*\*.
    - Data Completeness Met and Performance Met\*\* letter is represented in the Data Completeness and Performance Rate in the Sample Calculation listed at the end of this document. Letter a<sup>3</sup> equals 20 patients in the Sample Calculation.
  - b. If Patient identified as a tobacco user received tobacco cessation intervention (counseling and/or pharmacotherapy) equals No, proceed to check Documentation of medical reason(s) for not providing tobacco cessation intervention.
- 9. Check Documentation of medical reason(s) for not providing tobacco cessation intervention:
  - a. If Documentation of medical reason(s) for not providing tobacco cessation intervention equals Yes, include in Data Completeness Met and Denominator Exception.
    - Data Completeness Met and Denominator Exception letter is represented in the Data Completeness and Performance Rate in the Sample Calculation listed at the end of this document. Letter b<sup>2</sup> equals 10 patients in the Sample Calculation.
  - b. If Documentation of medical reason(s) for not providing tobacco cessation intervention equals No, proceed to check Patient identified as tobacco user did not receive tobacco cessation intervention (counseling and/or pharmacotherapy), reason not given.
- 10. Check Patient identified as tobacco user did not receive tobacco cessation intervention (counseling and/or pharmacotherapy), reason not given:
  - a. If Patient identified as tobacco user did not receive tobacco cessation intervention (counseling and/or pharmacotherapy), reason not given equals Yes, include in the Data Completeness Met and Performance Not Met.
    - Data Completeness Met and Performance Not Met letter is represented in the Data
       Completeness in the Sample Calculation listed at the end of this document. Letter c² equals 10 patients in the Sample Calculation.
  - b. If Patient identified as tobacco user did not receive tobacco cessation intervention (counseling and/or pharmacotherapy), reason not given equals No, proceed to check Data Completeness Not Met.
- 11. Check Data Completeness Not Met:
  - a. If *Data Completeness Not Met*, the Quality Data Code or equivalent was not submitted. 10 patients have been subtracted from the Data Completeness Numerator in the Sample Calculation.

#### Sample Calculations: Submission Criteria Two

Data Completeness equals Performance Met ( $a^3$  equals 20 patients) plus Denominator Exception ( $b^2$  equals 10 patients) plus Performance Not Met ( $c^2$  equals 10 patients) divided by Eligible Population/Denominator ( $d^2$  equals 50 patients). All equals 40 patients divided by 50 patients. All equals 80.00 percent.

Performance Rate equals Performance Met ( a<sup>3</sup> equals 20 patients) divided by Data Completeness Numerator (40 patients) minus Denominator Exception ( b<sup>2</sup> equals 10 patients). All equals 20 patients divided by 30 patients. All equals 66.67 percent.

\*See the posted measure specification for specific coding and instructions to submit this measure.

\*\*This measure defines tobacco cessation counseling as lasting 3 minutes or less. Services typically provided under CPT codes 99406 and 99407 satisfy the requirement of tobacco cessation intervention, as these services provide tobacco cessation counseling for 3-10 minutes. If a patient received these types of services, submit G-code G9906.

NOTE: Submission Frequency: Patient-Process

The measure diagrams were developed by CMS as a supplemental resource to be used in conjunction with the measure specifications. They should not be used alone or as a substitution for the measure specification.

#### **Submission Criteria Three:**

- Start with Denominator
- 2. Check Patient aged greater than or equal to 18 years:
  - a. If *Patient aged greater than or equal to 18 years* equals No, do not include in *Eligible Population/Denominator*. Stop processing.
  - b. If Patient aged greater than or equal to 18 years equals Yes, proceed to check At least two patient encounters during the performance period as listed in Denominator\*.
- Check At least two patient encounters during the performance period as listed in Denominator\*:
  - a. If At least two patient encounters during the performance period as listed in Denominator\* equals No, proceed to check At least one preventive encounter during the performance period as listed in Denominator\*.
  - b. If At least two patient encounters during the performance period as listed in Denominator\* equals Yes, include in Eligible Population/Denominator.
- 4. Check At least one preventive encounter during the performance period as listed in Denominator\*:
  - a. If At least one preventive encounter during the performance period as listed in Denominator\* equals No, do not include in Eligible Population/Denominator. Stop processing.
  - b. If At least one preventive encounter during the performance period as listed in Denominator\* equals Yes, include in Eligible Population/Denominator.
- 5. Denominator Population:
  - a. Denominator Population is all Eligible Patients in the Denominator. Denominator is represented as Denominator in the Sample Calculation listed at the end of this document. Letter d³ equals 100 patients in the Sample Calculation.
- 6. Start Numerator
- 7. Check Patient screened for tobacco use and received tobacco cessation intervention (counseling, pharmacotherapy, or both), if identified as a tobacco user:

- a. If Patient screened for tobacco use and received tobacco cessation intervention (counseling, pharmacotherapy, or both), if identified as a tobacco user equals Yes, include in Data Completeness Met and Performance Met\*\*.
  - Data Completeness Met and Performance Met\*\* letter is represented in the Data Completeness
    and Performance Rate in the Sample Calculation listed at the end of this document. Letter a<sup>4</sup>
    equals 20 patients in the Sample Calculation.
- b. If Patient screened for tobacco use and received tobacco cessation intervention (counseling, pharmacotherapy, or both), if identified as a tobacco user equals No, proceed to check Current tobacco non-user.
- 8. Check Current tobacco non-user.
  - a. If Current tobacco non-user equals Yes, include in Data Completeness Met and Performance Met.
    - Data Completeness Met and Performance Met letter is represented in the Data Completeness and Performance Rate in the Sample Calculation listed at the end of this document. Letter a<sup>5</sup> equals 20 patients in the Sample Calculation.
  - b. If Current tobacco non-user equals No, proceed to check Documentation of medical reason(s) for not screening for tobacco use.
- 9. Check Documentation of medical reason(s) for not screening for tobacco use:
  - a. If Documentation of medical reason(s) for not screening for tobacco use equals Yes, include in Data Completeness Met and Denominator Exception.
    - Data Completeness Met and Denominator Exception letter is represented in the Data Completeness and Performance Rate in the Sample Calculation listed at the end of this document. Letter b<sup>3</sup> equals 10 patients in the Sample Calculation.
  - b. If Documentation of medical reason(s) for not screening for tobacco use equals No, proceed to check Documentation of medical reason(s) for not providing tobacco cessation intervention if identified as a tobacco user.
- 10. Check Documentation of medical reason(s) for not providing tobacco cessation intervention if identified as a tobacco user:
  - a. If Documentation of medical reason(s) for not providing tobacco cessation intervention if identified as a tobacco user equals Yes, include in Data Completeness Met and Denominator Exception.
    - Data Completeness Met and Denominator Exception letter is represented in the Data Completeness and Performance Rate in the Sample Calculation listed at the end of this document. Letter b<sup>4</sup> equals 10 patients in the Sample Calculation.
  - b. If Documentation of medical reason(s) for not providing tobacco cessation intervention if identified as a tobacco user equals No, proceed to check Tobacco screening not performed OR tobacco cessation intervention not provided, reason not otherwise specified.
- 11. Check Tobacco screening not performed OR tobacco cessation intervention not provided, reason not otherwise specified:
  - a. If Tobacco screening not performed OR tobacco cessation intervention not provided, reason not otherwise

specified equals Yes, include in the Data Completeness Met and Performance Not Met\*\*\*.

- Data Completeness Met and Performance Not Met\*\*\* letter is represented in the Data
   Completeness in the Sample Calculation listed at the end of this document. Letter c³ equals 20 patients in the Sample Calculation.
- b. If Tobacco screening not performed OR tobacco cessation intervention not provided, reason not otherwise specified No, proceed to check Data Completeness Not Met.
- 12. Check Data Completeness Not Met:
  - a. If *Data Completeness Not Met*, the Quality Data Code or equivalent was not submitted. 20 patients have been subtracted from the Data Completeness Numerator in the Sample Calculation.

#### Sample Calculations: Submission Criteria Three

Data Completeness equals Performance Met ( a<sup>4</sup> plus a<sup>5</sup> equals 40 patients) plus Denominator Exception ( b<sup>3</sup> plus b<sup>4</sup> equals 20 patients) plus Performance Not Met ( c<sup>3</sup> equals 20 patients) divided by Eligible Population/Denominator ( d<sup>3</sup> equals 100 patients). All equals 80 patients divided by 100 patients. All equals 80.00 percent.

Performance Rate equals Performance Met ( a<sup>4</sup> plus a<sup>5</sup> equals 40 patients) divided by Data Completeness Numerator (80 patients) minus Denominator Exception ( b<sup>3</sup> plus b<sup>4</sup> equals 20 patients). All equals 40 patients divided by 60 patients. All equals 66.67 percent.

NOTE: Submission Frequency: Patient-Process

The measure diagrams were developed by CMS as a supplemental resource to be used in conjunction with the measure specifications. They should not be used alone or as a substitution for the measure specification.

<sup>\*</sup>See the posted measure specification for specific coding and instructions to submit this measure.

<sup>\*\*</sup>This measure defines tobacco cessation counseling as lasting 3 minutes or less. Services typically provided under CPT codes 99406 and 99407 satisfy the requirement of tobacco cessation intervention, as these services provide tobacco cessation counseling for 3-10 minutes. If a patient received these types of services, submit 4004F.

<sup>\*\*\*</sup>In the event that a patient is identified as a user but did not receive tobacco cessation intervention submit 4004F with 8P.